

Dejac

Dapoxetine Hydrochloride

Presentation

AC 30 Tablet: Each film coated tablet contains Dapoxetine Hydrochloride INN equivalent to Dapoxetine 30 mg.

Description

Dapoxetine is a selective serotonin reuptake inhibitor. Dapoxetine in premature ejaculation is presumed to be linked to the inhibition of neuronal reuptake of serotonin and the subsequent potentiation of the neurotransmitters action at pre and post synaptic receptors.

Dapoxetine is rapidly absorbed with maximum plasma concentrations (C_{max}) occurring approximately 1-2 hours after tablet intake. Greater than 99% of Dapoxetine is bound to human proteins and the metabolites of Dapoxetine are primarily eliminated in urine conjugates.

Indication

Dapoxetine Hydrochloride is indicated for the treatment of premature ejaculation in men.

Usage and Administration

recommended dose of Dapoxetine Hydrochloride (Dejac) is as follows -

Patients	Dosage	
Adult Men to 64 years of age)	30 mg	Once Daily 1 to 3 hours prior to sexual activity
Dose may be increased to the maximum recommended dose of 60 mg.		
Children & adolescents	Dapoxetine should not be used in individuals below 18 years of age.	

Contraindication

Dapoxetine Hydrochloride is contraindicated in patients with significant pathological cardiac conditions (Heart failure, conduction abnormalities, significant Ischemic Heart Disease, concomitant treatment with Monoamine Oxidase Inhibitors, Thioridazine, Selective Serotonin Reuptake Inhibitors) & history of hypersensitivity to any component of the product.

Caution

Patients should be advised not to use Dapoxetine in combination with recreational drugs. Recreational drugs with serotonergic activity such as ketamine, methylenedioxymethamphetamine (MDMA) and lysergic acid diethylamide (LSD) may lead to potentially serious reactions if combined with Dapoxetine. These reaction includes - tachycardia, hyperthermia and serotonin syndrome. Use of Dapoxetine with recreational drugs with sedative properties such as narcotics and benzodiazepines further increase somnolence and dizziness.

Effect

The most common side effects are nausea, dizziness, dry mouth, headache, diarrhea and insomnia.

Pregnancy & lactation

Dapoxetine is not indicated for use in women.

Drug Interaction

Potential serious reactions with MAOIs. Inhibits metabolism of Thioridazine. Risk of serotonin associated effects with serotonergic medicinal/herbal products (Lysergic acid, triptans, Tramadol, Linezolid, SSRIs, SNRIs, Lithium & St. John's wort). CNS active medicinal products. Possible reduction of clearance with CYP2D6, CYP3A4 & flavin monooxygenase 1 inhibitors.

Overdose

Overdose with Dapoxetine Hydrochloride has been rare. In general, symptoms of overdose with SSRIs include serotonin-mediated adverse reactions such as somnolence, gastrointestinal disturbance (such as nausea and vomiting), tachycardia, hypertension, agitation and dizziness.

Storage

Store in a cool & dry place, away from light and Keep out of reach of the children.

Commercial Pack

AC 30 Tablet: Each box contains 1 x 4 tablets in blister strip.

Manufactured by:
GLOBE PHARMACEUTICALS LTD.
Bangladesh

ডিজেক

ড্যাপোরেটিন হাইড্রোচ্লোরাইড

উপস্থাপন

ডিজেক 30 ট্যাবলেট : c^oZU idj f KtUW U'vetj tu i tqQ W'tcvf. Ub nBtWtKv vBW AitGcb
W'tcvf. Ub 30 mgM Gi mgZj |

বিবরণ

W'tcvf. Ub GKU mtj wf tmf vUmb vi AvctUK BbmieUi | W'tcvf. Ub mgZ tmf vU
cping MhYK evamf f Kivi gta'g mGes tcvo mBtbP K vi tmPj ,tj vi KvhKwi ZtK eU
W'tcvf. Ub tmfbi 1-2 Njvi gta' covRgtZ mteP givvq tcstQ| 99% Gi tekx W'tcv
gibe t'ni i t'i tcUtb Ave x nq Ges W'tcvf. Ub tgUtev vBUm c^ogKfite chtei
ib:mZ nq |

নির্দেশনা

W'tcvf. Ub nBtWtKv vBW c^oBeq c^oej af i Dch mgfqi c^oeB exHrtZi vPKrmiq ibt R
সেবনমাত্রা ও প্রয়োগবিধি
ubgij LZ givvq W'tcvf. Ub nBtWtKv vBW (ডিজেক) ibt RkZ -

রোগী	মাত্রা	
প্রাঞ্চবয়ক (১৮ থেকে ৬৪ বছর)	30 mgM	%nBK GKevi mnevtmi 1 t_k 3 Njv cte
hi` 30 mgM Gi KvhKwi ZtSf vRbK bv nq Zte Gi givv 60 mgM chsf evotbv thtz c		
শিশু	18 e0ti i bP ti vMx t i t'j f W'tcvf. Ub eenvi Kiv DIPZ bq	

প্রতিনির্দেশনা

th mKj ti Mxi Dfj oLthM c^ovj lRKj KvhKwi mgm v (nU^cdbj y KbWkkb Gebig
Dfj oLthM BktKngK nV^cW^cRm, gtbvA^cg^cbv A^c fWR BbmieUi, _v^cqmi WmRb, w
tmf vUmb vi AvctUK BbmieUi) Ges W'tcvf. Ub nBtWtKv vBW Gi tKv Dcr vtb
AvZmste bkj Zv Avt Zt i t'j f W'tcvf. Ub c^oZibt RkZ |

সর্তরতা

W'tcvf. Ub bi mft w^cqkbj W^cmi eenvi t_k K wei Z vKv DIPZ | w^cqkbj WM th
WkUmgb, ig_Bj bWBAr^cg^cndUwgb I j Bm^cW^cRk GmW W^cBB_vBj GgvB^cWi (Gj GmW)
W'tcvf. Ub bi eenvi i^cz Zi c^oZipqf t Lv w^cz c^ovj | c^oZipqf gta' - Gvi
nBcvi vngq Ges tmf vUmb imbÜg ciuj ny Z| bvi KvhKwi Ges tebtRwqvwRvcb WM Gi
th ib^cffve ciuj ny Z nq, W'tcvf. Ub bi mft w^cqkbj WM e eUZ ntj Zv Avt vfe h
পার্শ্বপ্রতিক্রিয়া

cikf^clZipqf gta' itqfQ eng eng five, ib^cffve, gj^c i^cKq hv^c qv, gv ve v, Wqui qv Ges Aib

গর্ভবত্তায় ও দুর্ঋান্তকালে ব্যবহার

W'tcvf. Ub nBtWtKv vBW gnf v i t'j f W'tcvf. Ub c^oZibt RkZ bq |

ঔষধের আঙ্গক্রিয়া

W'tcvf. Ub bi mft Ggj AvBGm eenvi i^cz Zi c^oZipqf ciuj ny Z nq | _v^cqmi Wm
fgUewj Rg eU Ktj | tmf vUmb WM A^c fIR `te i (Gj -ut^ctUtdb, ut^cUvb, Ut
ij tbvRw W, GmGmAvi AvBGm, GmGmAvi AvBGm, ij^cqig Ges GmW, tRvb^cm I U^ctc^c.)
tmf vUmb vi Rk c^ofite SiK _vK | w^cGBGm I l^c Gi mft KvhKwi c^oZipqf t Lv
w^cl qBic2W6, w^cl qBic3G4 Ges dwfb gtbvA^c R^cbm 1 BbmieUi, JIai ib:miY Kigtq
মারোধিক্রিয়া

W'tcvf. Ub nBtWtKv vBW Gi t'j f givvafK i Z_ weij | mvavibfve AvZm
GmGmAvi AvB tmfbi dtj msNuz c^oZipqf i j y b mft thgb- Aib t, eng eng five, eng
`m b Gi ex, K^cub, Avt vob Ges ib^cffve t Lv w^cz c^ovj |

সংরক্ষণ

Avt v t_k K^c, t^c | W^cUv t^cb Ges w^cki t^c bimtj i evBt i vLp |

বাণিজ্যিক মোড়ক

ডিজেক 30 ট্যাবলেট : c^oZ evt i tqfQ 1 x 4 u U'vetj U Gi weo vi /c |



প্রস্তুতকারক:
গ্লোব ফার্মাসিউটিক্যালস লিঃ
eisj f k