

Syrup
ZIKID
Zinc sulphate USP

PRESENTATION :

ZIKID Syrup : Bottle containing 100 ml syrup. Each 5 ml Syrup contains Zinc sulphate monohydrate USP equivalent to elemental Zinc 10 mg.

INDICATION :

ZIKID Is indicated for the treatment of Recurrent respiratory tract infections, diarrhoea, loss of appetite, severe growth retardation, deformed bone formation, impaired immunological response, acrodermatitis enteropathica, parakeratotic skin lesions, defective and delayed wound healing, Anaemia, night blindness, mental disturbances.

DOSAGE & ADMINISTRATION :

Children under 10 kg : One teaspoonful 2 times daily.
Children within 10 to 30 kg : Two teaspoonfuls 1-3 times daily.
Adults and children over 30 kg body weight : Four teaspoonfuls 1-3 times daily.

CONTRA-INDICATION :

It is contraindicated in patients with hypersensitivity to Zinc.

PRECAUTION & WARNING :

Concurrent administration of Zinc salt with penicillamine might diminish the effect of penicillamine. The absorption of Zinc although poor, may be decreased by various compounds including some foods.

DRUG INTERACTIONS :

Zinc sulphate may inhibit the absorption of concurrently administered tetracycline, when both are being given an interval of at least 3 hours should be allowed.

OVERDOSE :

Zinc sulphate is corrosive in over dose. Symptoms are corrosion and inflammation of the mucous membrane of the mouth and stomach; ulceration of the stomach followed by perforation may occur. Gastric lavage and emesis should be avoided. Demulcents such as milk should be given. Chelating agents such as sodium edetate may be useful.

PHARMACEUTICAL PRECAUTION :

Store in a cool, dry place. The syrup should be protected from light.

HOW SUPPLIED :

Bottle containing 100 ml ZIKID Syrup.

Manufactured by:



Globe Pharmaceuticals Ltd.

Noakhali, Bangladesh.